



# Reach for the Sky

**A New Clinical Research Study for Children, Teens, and Adults  
Diagnosed With Lennox-Gastaut Syndrome – The Skyway Study**

## What Study Participants Can Expect




To learn more about this new study, or to find out if you, your child, or a loved one may qualify, please visit the Skyway Study website at [skyway-study.com](http://skyway-study.com) or [clinicaltrials.takeda.com](http://clinicaltrials.takeda.com).



People who are eligible for the study and decide to participate will receive:

- The investigational medication or placebo and all study-related care at no cost
- Care and close monitoring during the study from doctors and staff who understand their condition
- The option to join the open-label extension (OLE) study after completing the study where all participants will receive the investigational medication
- Reimbursement for expenses related to study participation such as travel, meals, etc. may be available

To help make an informed decision on whether or not to take part in this study, potential participants will go through a process called informed consent. The informed consent process will review what is involved in participating in the study and the potential risks and benefits. A participant can leave the study at any time, for any reason, without penalty.






## A Person May Qualify for the Skyway Study If He/She:

- Is 2 to 35 years old
- Has been diagnosed with Lennox-Gastaut syndrome
- Has 8 or more major motor drop (MMD) seizures each month for the last 3 months
- Has been unable to control seizures with at least 2 anti-seizure medications in the past

Study staff will determine eligibility based on additional study criteria.

## What Is Lennox-Gastaut Syndrome?

Lennox-Gastaut syndrome (LGS) is a rare epilepsy syndrome associated with severe seizures in childhood. Diagnosed during infancy or early childhood, LGS is characterized by uncontrollable movements of the body which are sometimes difficult to treat with currently approved anti-seizure medications.



## About the Skyway Study

The Skyway Study is a Phase 3 clinical research study of an investigational medication for children, teens, and adults diagnosed with Lennox-Gastaut syndrome who are currently using anti-seizure medications to manage symptoms and are still experiencing seizures. The investigational medication, soticlestat, works differently than currently approved medications and is being tested as an add-on therapy in the clinical research study.

The goal of the Phase 3 clinical research study is to assess the effectiveness, safety and tolerability of the investigational medication when taken with other anti-seizure treatments. In previous clinical research studies, the investigational drug has been shown to reduce the number of seizures in study participants.

During the study, participants will be split into two groups: one will receive the study drug and one will receive placebo (looks like the study drug but contains no active ingredients). After completing the study, participants can join the open-label extension (OLE) study where every participant will receive the study drug.

## Study Participation Involves:

- Up to 5 planned in-person study visits
- Approximately 25 weeks of study participation

## What Will Occur During the Study?

- **Screening (4–6 weeks):** During screening, it will be determined whether you, your child, or a loved one qualifies for the study.
- **Titration Period (4 weeks):** During the first visit of the four-week titration period, participants will be randomized into one of two groups. One group will receive the study drug, and one group will receive placebo. The placebo will look like the study drug but have no active ingredients. Titration is the process of adjusting a dose of medication to determine the dose a person can tolerate. During the four weeks, participants will receive the study drug or placebo at increased doses. The dose will be increased throughout the titration period as long as participants don't have any tolerability issues. If participants can't tolerate the minimum dose, they will discontinue from the study.
- **Maintenance Period (12 weeks):** The dose a participant receives at the end of the titration period will be taken throughout the 12-week maintenance period. All participants may have the option to join the open-label extension (OLE) study after completing the maintenance period (based on selection criteria for the OLE study). During the OLE study, all participants (including those who received placebo) will take the investigational drug.
- **Follow-up (3 weeks):** Participants who are not joining the OLE study will taper off the study drug over one week and complete a safety follow-up visit or phone call approximately two weeks after receiving their last dose of the study drug.

Throughout participation in the study, all study-related visits, tests, and study medication will be provided at no cost. Reimbursement for expenses related to study participation such as travel, meals, etc. may also be available.